



## Early Journal Content on JSTOR, Free to Anyone in the World

This article is one of nearly 500,000 scholarly works digitized and made freely available to everyone in the world by JSTOR.

Known as the Early Journal Content, this set of works include research articles, news, letters, and other writings published in more than 200 of the oldest leading academic journals. The works date from the mid-seventeenth to the early twentieth centuries.

We encourage people to read and share the Early Journal Content openly and to tell others that this resource exists. People may post this content online or redistribute in any way for non-commercial purposes.

Read more about Early Journal Content at <http://about.jstor.org/participate-jstor/individuals/early-journal-content>.

JSTOR is a digital library of academic journals, books, and primary source objects. JSTOR helps people discover, use, and build upon a wide range of content through a powerful research and teaching platform, and preserves this content for future generations. JSTOR is part of ITHAKA, a not-for-profit organization that also includes Ithaka S+R and Portico. For more information about JSTOR, please contact [support@jstor.org](mailto:support@jstor.org).

*SOLUTION OF THE GENERAL CASE OF PROBLEM. 87,  
MATH. VISITOR, NO. 2.*

BY PROF. WM. WOOLSEY JOHNSON, ANNAPOLIS, MARYLAND.

LET there be a series of casks containing  $a_1, a_2$  etc. gallons, of which the first is filled with a mixture containing  $a$  gallons of wine, and the rest are filled with water. Water runs into the first cask at the rate of  $m$  gallons per minute, the mixture escapes into the second cask at the same rate, and the mixture in the second cask escapes into the third at the same rate, and so on. It is required to find the quantity of wine in each cask at the end of  $t$  minutes, the fluids being supposed to mingle perfectly.

Denoting the required quantities by  $x_1, x_2$  etc., my published solution of the special case (Math. Visitor, Jan., 1879) suggests what will be verified in the following solution; namely, that the quantities take the form,

$$\left. \begin{aligned} x_1 &= A_{1,1} e^{-\frac{mt}{a_1}} \\ x_2 &= A_{1,2} e^{-\frac{mt}{a_1}} + A_{2,2} e^{-\frac{mt}{a_2}} \\ x_3 &= A_{1,3} e^{-\frac{mt}{a_1}} + A_{2,3} e^{-\frac{mt}{a_2}} + A_{3,3} e^{-\frac{mt}{a_3}} \\ &\vdots \\ x_n &= A_{1,n} e^{-\frac{mt}{a_1}} + A_{2,n} e^{-\frac{mt}{a_2}} + \dots + A_{n,n} e^{-\frac{mt}{a_n}} \end{aligned} \right\} \quad (1)$$

in which  $A_{1,1}$  etc., are constants to be determined. The first subscript refers to the exponential to which the coefficient belongs and the second to the variable in whose expression it occurs. It is to be remarked that  $A_{r,n}$  vanishes when  $r > n$ .

Since  $x_1 \div a_1$  represents the fractional part of the mixture in the first cask which consists of wine,  $m(x_1 \div a_1)$ ,  $m(x_2 \div a_2)$ , etc., denote the rates at which wine is escaping from the 1st, 2nd, etc. casks, and, at the same time, the rate at which wine is entering the 2nd, 3rd, etc. casks. Hence the conditions of the problem give the differential equations,

$$\left. \begin{aligned} \frac{dx_1}{dt} &= \quad \quad \quad - \frac{mx_1}{a_1} \\ \frac{dx_2}{dt} &= \frac{mx_1}{a_1} - \frac{mx_2}{a_2} \\ \frac{dx_3}{dt} &= \frac{mx_2}{a_2} - \frac{mx_3}{a_3} \\ &\vdots \\ \frac{dx_n}{dt} &= \frac{mx_{n-1}}{a_{n-1}} - \frac{mx_n}{a_n} \end{aligned} \right\}, \quad (2)$$

and, putting  $t=0$  in (1), we have, to determine the constants of integration,



may be determined. Thus from (3) we have  $A_{1,1} = a$ , hence from (6)

$$A_{1,n} = \frac{a_1^{n-2} a_n a}{(a_1 - a_2)(a_1 - a_3) \dots (a_1 - a_n)}. \quad (7)$$

Again, making  $n = 2$  in (7),

$$A_{1,2} = \frac{a_2 a}{a_1 - a_2},$$

substituting in the second of group (3), we derive

$$A_{2,2} = \frac{a_2 a}{a_2 - a_1},$$

whence, from (6),

$$A_{2,n} = \frac{a_2^{n-2} a_n a}{(a_2 - a_1)(a_2 - a_3) \dots (a_2 - a_n)}. \quad (8)$$

The form of equations (7) and (8) suggests that in general

$$A_{r,n} = \frac{a_r^{n-2} a_n a}{(a_r - a_1)(a_r - a_2) \dots (a_r - a_n)} \quad (9)$$

(the subscripts of the second letter in the binomials taking all values from 1 to  $n$  except  $r$ ), and this expression will be demonstrated when it is shown that it satisfies the  $n$ th equation of group (3); that is, the equation

$$A_{1,n} + A_{2,n} + A_{3,n} + \dots + A_{n,n} = 0.$$

Making the substitutions, dividing by  $a_n a$ , and multiplying by the expression

$$(a_1 - a_2)(a_1 - a_3) \dots (a_1 - a_n) \times (a_2 - a_3)(a_2 - a_4) \dots (a_2 - a_n) \\ \times (a_3 - a_4) \dots (a_{n-1} - a_n),$$

the equation to be verified becomes

$$\left. \begin{aligned} &+ a_1^{n-2} (a_2 - a_3) \dots (a_2 - a_n) \cdot (a_3 - a_4) \dots (a_3 - a_n) \dots (a_{n-1} - a_n) \\ &- a_2^{n-2} (a_1 - a_3)(a_1 - a_4) \dots (a_1 - a_n) \cdot (a_3 - a_4) \dots (a_{n-1} - a_n) \\ &+ a_3^{n-2} (a_1 - a_2)(a_1 - a_4) \dots (a_1 - a_n) \dots \\ &\dots \dots \dots \end{aligned} \right\} = 0,$$

in which the coefficient of  $a_r^{n-1}$  is the product of all the differences which do not involve  $a_r$ . Using the determinant expressions for these products, the equation takes the form

$$a_1^{n-2} \begin{vmatrix} 1, & 1 & \dots & 1 \\ a_n, & a_{n-1}, & \dots & a_2 \\ a_n^2, & a_{n-1}^2, & \dots & a_2^2 \\ \vdots & \vdots & & \vdots \\ a_n^{n-2}, & \dots & \dots & a_2^{n-2} \end{vmatrix} - a_2^{n-2} \begin{vmatrix} 1, & 1, & \dots, & 1, & 1 \\ a_n, & \dots & a_3, & a_1 \\ a_n^2, & \dots & a_3^2, & a_1^2 \\ \vdots & & \vdots & \\ \vdots & & \vdots & \\ a_n^{n-2}, & \dots & a_1^{n-2} \end{vmatrix} + a_3^{n-2} \begin{vmatrix} 1; & 1 \dots \\ a_n & \dots \\ a_n^2 & \dots \text{etc.} \\ \vdots & \\ a_n^{n-2} & \dots \end{vmatrix} = 0.$$

Now the right hand member of this equation is equivalent to the determinant

$$\begin{vmatrix} 1 & , & 1 & , & \dots & 1 & , & 1 \\ a_n & , & a_{n-1} & , & \dots & a_2 & , & a_1 \\ a_n^2 & , & a_{n-1}^2 & , & \dots & a_2^2 & , & a_1^2 \\ \vdots & & \vdots & & & \vdots & & \vdots \\ \vdots & & \vdots & & & \vdots & & \vdots \\ \vdots & & \vdots & & & \vdots & & \vdots \\ a_n^{n-2} & , & a_{n-1}^{n-2} & , & \dots & a_2^{n-2} & , & a_1^{n-2} \\ a_n^{n-2} & , & a_{n-1}^{n-2} & , & \dots & a_2^{n-2} & , & a_1^{n-2} \end{vmatrix}$$

which is identically = 0, because the last two rows are identical. Hence the solution of the problem is given by (1), the coefficients being determined by (9).

It is a remarkable result of this solution that the value of the fraction  $x_n \div a_n$ , or ratio of the wine to the whole contents of the last cask, is wholly independent of the order of the casks.

---

### NOTE ON A FORM OF THE EQ. OF THE TANG. TO A CONIC.

---

BY PROF. W. W. HENDRICKSON, U. S. NAVAL ACADEMY.

THE envelope of any line  $A\alpha^2 + B\alpha + C = 0$ , where  $\alpha$  is an arbitrary parameter, and  $A, B$  and  $C$  are functions of  $x$  and  $y$ , is  $B^2 = 4AC$ ; it follows that if the equation to any conic be put in the form  $B^2 = 4AC$ , the equation to the tangent may be written either as  $A\alpha^2 + B\alpha + C = 0$  or  $C\alpha^2 + B\alpha + A = 0$ .

The equations to the conic sections and their tangents may be written as follows:

Circle,  $y^2 = (a+x)(a-x)$ , tangent,  $(a-x)\alpha^2 + 2ya + a + x = 0$ .

Parabola,  $y^2 = 4ax$ , "  $a\alpha^2 + ya + x = 0$ .

Ellipse,  $a^2y^2 = b^2(a+x)(a-x)$ , "  $b(a-x)\alpha^2 + 2aya + b(a+x) = 0$ .

Hyperb.,  $a^2b^2 = (bx+ay)(bx-ay)$ , "  $(bx-ay)\alpha^2 + 2aba + bx + ay = 0$ .

This form of the equation to the tangent will be found useful in certain problems of Analytical Geometry; thus, suppose it required to find the tangents to the conic

$$y^2 - 2xy + 2x^2 + 8y + 6x + 49 = 0,$$

which pass through  $(-14, -22)$ . The equation may be put in the form  $(y - x + 4)^2 + (x + 11)(x + 3) = 0$ , and the equation to the tangent is  $(x + 11)\alpha^2 + 2(y - x + 4)\alpha - x - 3 = 0$ ; substituting the co-ordinates of the given point, we have  $3\alpha^2 + 8\alpha - 11 = 0$ , whence  $\alpha = 1$  or  $-\frac{11}{3}$ , and the equations to the required tangents are  $y - x + 8 = 0$ , and  $89x - 33y + 520 = 0$ .